

令和7年度 フェロモントラップ調査結果(コナガ)







掲載元(「ひろしま病害虫情報」で検索)

https://www.pref.hiroshima.lg.jp/site/byogaichu/

調査データ

調査データ							
地带区分		中東部			北西部		
設置場所		世羅郡世羅町下津田			山県郡北広島町南門原		
周辺作物		キャベツ			キャベツ		
月	半旬	本年	平均(1年)	前年	本年	平年	前年
5月	1	9.3	6.7	6.7	96.9	7.2	7.5
	2	21.7	4.2	4.2	88.4	9.9	6.2
	3	24.9	6.1	6.1	75.7	12.8	4.3
	4	4.3	10.0	10.0	48.1	10.8	2.9
	5	8.2	6.4	6.4	43.2	9.3	6.3
	6	8.7	4.7	4.7	33.0	10.4	7.7
6月	1	9.4	3.3	3.3	10.7	4.8	2.1
	2	22.1	2.9	2.9	5.0	5.5	11.4
	3	7.8	7.1	7.1	50.9	6.7	6.3
	4	1.7	3.3	3.3	70.9	5.1	3.9
	5	0	0.7	0.7	28.6	6.0	7.9
	6	7.9	0.7	0.7	16.6	5.6	5.0
7月	1	4.1	0.7	0.7	13.9	4.2	3.7
	2	2.0	1.6	1.6	14.3	3.8	2.9
	3	2.5	2.9	2.9		5.1	0.7
	4		0.7	0.7		4.6	0.7
	5		1.6	1.6		2.7	0.6
	6		2.0	2.0		3.0	0
8月	1		0.6	0.6		3.4	1.4
	2		0.8	0.8		3.5	1.6
	3		0.3	0.3		5.9	1.7
	4		0	0		14.8	11.8
	5		0	0		17.4	20.6
	6		0	0		13.5	36.0
9月	1		2.0	2.0		11.2	13.7
	2		2.0	2.0		8.9	2.9
	3		0	0		8.6	2.9
	4		0.6	0.6		6.4	8.6
	5		0.4	0.4		10.1	6.9
	6		0	0		8.2	2.2
10月	1		0	0		7.8	1.5
	2		0	0		7.5	1.3
	3		0	0		7.1	1.7
	4		0	0		4.5	2.9
	5		0	0		7.5	0.6
	6		0	0		8.6	0