



「広島県病害虫発生予察調査データ」

令和7年度 フェロモントラップ等調査結果(南東部・南西部)

ハナアザミウマ (青色粘着板)

トラップ調査結果

(1) グラフ



(2)調査データ

地帯区分		南東部			南西部		
設置場所		尾道市瀬戸田町B			呉市倉橋町		
周辺作物		かんきつ			かんきつ		
月	半旬	本年	平均(4)	前年	本年	平均(3)	前年
5月	1	3.6	21.0	6.0	0	17.4	2.9
	2	6.0	27.3	33.4	4.9	13.7	5.4
	3	6.0	35.2	68.6	8.1	7.1	6.3
	4	6.0	64.1	52.9	43.1	18.1	2.9
	5	11.8	105.2	26.7	43.1	50.8	14.9
	6	17.3	138.6	15.7	51.8	95.9	24.4
6月	1	8.5	40.5	20.7	32.0	96.4	21.4
	2	8.5	40.5	10.8	32.0	64.0	15.7
	3	15.8	32.7	14.2	32.0	39.4	13.4
	4	20.7	44.2	44.0	49.3	33.6	35.1
	5	20.7	187.6	160.0	49.3	56.0	68.6
	6	113.3	474.1	189.7	49.3	96.1	65.7
7月	1	136.4	594.1	155.1	36.6	95.5	34.4
	2	136.4	362.5	92.1	36.6	92.5	27.4
	3	23.9	131.3	74.3	36.6	105.7	82.9
	4	23.9	82.3	56.3	39.6	123.4	93.1
	5	22.8	73.0	39.0	40.3	117.4	77.1
	6	21.9	36.4	21.4	48.4	79.1	54.9
8月	1	18.2	10.6	3.6	8.7	27.4	27.9
	2	10.9	5.9	2.7	8.7	12.7	13.0
	3	6.1	4.4	2.1	8.7	4.6	7.6
	4	6.1	3.4	2.1	0	4.3	5.0
	5	2.1	3.1	2.7	0	3.2	5.7
	6	1.3	2.5	2.6	0	3.0	5.7
9月	1	0.9	1.0	1.1	0.3	2.0	2.9
	2	0	2.3	0	0.3	1.9	0.7
	3	0	12.1	1.0	0.3	1.9	0
	4	9.6	19.5	3.6	7.3	18.5	16.7
	5	23.9	19.3	6.4	7.3	43.9	43.0
	6	23.9	45.1	13.6	7.3	96.4	103.6
10月	1	47.4	43.5	12.7	37.3	96.7	134.7
	2	63.0	32.1	12.4	37.3	82.5	127.6
	3	63.0	110.9	13.6	37.3	42.4	67.9
	4	26.1	271.5	24.4	11.9	45.8	73.0
	5	1.4	354.1	37.3	11.9	40.0	73.2
	6	1.7	381.9	63.0	14.3	67.6	181.3