




令和8年度 フェロモントラップ調査結果(ハスモンヨトウ)

※調査データはグラフの下にあります。






調査データ

地帯区分		中西部			北西部			北西部		
設置場所		東広島市八本松町・農業技術C			北広島町南門原			北広島町木次		
周辺作物					キャベツ			キャベツ		
月	半旬	本年	平年	前年	本年	平年	前年	本年	平年	前年
5月	1	1.8	0.4	0	0	0.1	0	3.4	0.2	0
	2	1.8	0.8	0	0.4	0.4	0	2.6	0.7	1.0
	3	2.0	5.7	0	1.2	1.2	0	1.0	1.5	0
	4	0.4	8.2	5.6	3.1	4.4	1.9	5.0	1.3	0
	5	4.8	10.9	7.0	16.3	3.7	2.8	0.8	3.4	0
	6	3.7	9.8	6.4	24.3	3.0	3.3	0.2	2.0	3.3
6月	1	7.7	8.2	5.0	10.2	1.9	0	1.7	1.0	2.7
	2	10.6	11.3	4.2	7.5	2.3	0	1.3	2.4	3.5
	3	5.4	21.1	19.2	6.5	5.4	7.7	0	4.8	29.1
	4	7.2	29.5	21.2	24.3	14.2	19.4	0.3	10.8	66.4
	5	16.8	28.8	17.1	51.9	33.8	45.7	1.7	30.9	86.7
	6	4.6	19.7	7.9	16.4	32.5	29.1	2.5	28.6	54.1
7月	1	0.2	23.9	3.4	26.1	32.9	25.3	9.2	40.7	17.2
	2	1.6	24.7	1.8	34.1	25.0	25.7	8.8	58.3	12.4
	3	5.2	21.3	4.3	16.4	23.5	6.4	14.2	51.2	12.0
	4	3.6	17.7	1.7	44.3	23.6	6.4	24.3	28.4	9.9
	5	5.4	14.5	0.3	56.4	30.9	13.3	0	25.4	6.7
	6	15.4	18.4	3.3	92.3	44.7	48.9	18.9	38.2	19.7
8月	1	19.6	19.5	4.7	76.9	44.1	68.6	27.1	54.7	153.3
	2	23.4	33.7	8.0	115.4	70.0	42.8	35.5	115.9	131.5
	3	16.8	37.2	6.1	125.0	104.9	25.6	50.8	150.7	80.8
	4	27.0	34.2	3.9	125.0	117.8	38.8	60.0	143.4	66.3
	5	27.6	37.1	97.9	127.7	149.5	87.5	26.7	128.7	107.8
	6	42.1	88.0	125.1	155.5	270.3	204.3	179.1	189.8	164.5
9月	1	79.3	62.4	14.9	132.2	220.7	250.0	131.2	168.8	263.8
	2	55.6	58.9	10.0	133.9	150.2	128.5	109.7	163.4	204.0
	3	71.0	50.0	5.8	133.9	119.2	47.5	94.7	149.8	147.2
	4	35.0	49.6	10.6	48.6	143.4	405.7	76.8	194.8	421.4
	5	24.5	53.1	11.2	27.2	124.5	211.8	27.5	184.7	425.0
	6	61.7	42.3	11.0		81.2	76.1	72.5	114.8	300.0
10月	1		46.5	16.6		64.6	66.4		133.2	275.0
	2		64.7	39.6		38.6	3.9		104.7	201.7
	3		54.9	27.4		20.2	13.5		45.6	131.7
	4		59.1	19.8		21.3	52.1		67.7	191.7
	5		67.3	19.0		30.6	15.6		47.4	35.8
	6		64.7	7.1		23.9	58.0		56.3	133.3
11月	1		3.1	3.1		0	4		2.1	29.1
	2		3.8	7.1		1.0	1.0		1.0	63.3
	3		0.8	10.0		1.9	1.9		3.6	30.0
	4		0	16.4		2.1	2.1		2.1	18.3
	5		0	0		0	0		0.4	3.7
	6		3.0	3.0		0	0		0	9.0

令和7年度 フェロモントラップ調査結果(ハスモンヨトウ)

※調査データはグラフの下にあります。



調査データ

地帯区分		南東部			中東部			中東部		
設置場所		福山市新市町			世羅町山中福田			三次市廻神町		
周辺作物		アスパラガス			ねぎ			だいず		
月	半旬	本年	平年	前年	本年	平均(9年)	前年	本年	平均(1年)	前年
5月	1	0	0.4	0	1.1	1.4	0	0	0.8	1
	2	0	0.6	0.9	1.1	2.1	0	1.0	1.0	1
	3	0	0.8	2.1	1.0	4.0	0	2.0	1.8	2
	4	0	2.9	7.1	0.7	7.7	0	3.0	7.6	8
	5	0.9	3.6	6.7	13.9	16.7	0	5.0	17.8	18
	6	1.6	4.0	6.6	39.0	14.3	1.1	36.0	23.0	23
6月	1	2.1	3.6	7.1	40.3	8.2	4.5	99.0	10.0	10
	2	10.7	4.5	6.4	10.0	7.6	9.0	57.5	4.0	4
	3	6.9	6.7	15.9	7.1	11.5	26.4	49.5	25.0	25
	4	4.6	9.2	23.0	5.6	23.1	72.4	25.0	87.0	87
	5	5.7	13.0	26.4	4.3	28.6	76.7	175.0	66.0	66
	6	5.1	14.8	24.1	75.0	32.6	40.7	73.7	84.0	84
7月	1	9.0	17.6	33.9	42.0	42.8	19.6	55.3	72.0	72
	2	15.0	21.1	49.3	16.3	50.2	30.6	66.0	179.0	179
	3	7.9	22.9	63.6	1.4	57.1	55.0	12.0	102.5	103
	4	6.6	26.1	46.9	44.3	43.5	25.0	26.0	50.5	51
	5	7.9	29.8	34.4	55.7	46.4	32.3	23.3	35.0	35
	6	19.7	43.1	35.3	72.4	56.6	51.8	27.0	66.0	66
8月	1	14.3	53.2	55.2	83.6	61.9	116.7	16.7	100.0	100
	2	22.0	65.8	90.2	87.4	85.5	87.9	18.0	219.2	219
	3	25.1	74.9	97.9	97.9	138.8	59.1	35.0	175.8	176
	4	17.1	79.8	52.1	98.4	169.6	38.8	34.0	143.7	66
	5	50.9	102.5	97.9	75.0	160.8	87.5	33.6	107.8	108
	6	60.4	158.1	125.1	164.0	231.8	204.3	87.4	164.5	165
9月	1	40.0	157.1	102.3	216.1	165.2	150.3	68.0	196.0	264
	2	34.3	164.3	114.3	217.6	157.4	109.4	23.0	194.0	194
	3	21.1	133.3	64.6	93.6	172.5	190.7	20.0	195.0	195
	4	23.6	128.2	70.4	169.6	182.8	216.9	33.0	290.0	290
	5	32.1	132.2	97.9	217.1	171.7	237.4	10.8	255.0	255
	6		84.6	59.3	260.0	122.0	250.0		384.6	385
10月	1		103.7	70.6		134.3	219.6		399.4	399
	2		109.2	78.1		132.6	138.7		423.0	423
	3		75.7	35.0		109.0	85.6		506.0	506
	4		74.1	88.7		108.0	149.7		470.0	470
	5		63.9	83.0		100.3	122.3		346.0	346
	6		54.9	7.1		85.1	58.0		133.3	133
11月	1		9.3	3.1		4.3	4		29.1	29
	2		7.0	7.1		10.2	1.0		63.3	63
	3		7.9	10.0		7.2	7.2		30.0	30
	4		10.0	16.4		3.9	3.9		18.3	18
	5		0	0		2.1	2.1		3.7	4
	6		0	0		0.4	0.4		9.0	9